

Optimization of ELM control using RMPs with minimal fast-ion loss


Youwen Sun¹, Yaoning Zhang^{1,2}, Kaiyang He¹, Xuemin Wu^{1,2}, Yueqiang Liu³

¹ Institute of Plasma Physics, Chinese Academy of Sciences, Hefei 230031, China

² University of Science and Technology of China, Hefei 230026, China


³ General Atomics, PO Box 85608, San Diego, CA 921865608, USA

Email: ywsun@ipp.ac.cn





The 16th International West Lake Symposium on
the “Physics of Fusion and Space Plasmas” Apr. 10 - 12 2026, Hangzhou, China






ITER, $n = 3, 4$






EAST, $n = 1-4$

- RMP ELM control required in ITER
- EAST RMP covers the same toroidal mode number up to $n=4$

Key issue for understanding 3D physics





Vacuum Plasma response to RMP Response



- Resonant RMP on ELM suppression
- Non-resonant RMP on fast-ion loss
- Optimized RMP ELM suppression with minimal fast-ion loss

ELM suppression achieved by $n=4$ RMPs in ITER-like low torque plasmas in EAST





[Sun Y. *et al.*, *NF* 61, 106037 (2021)]

- Experimental condition is close to low torque ITER type-I ELMy H-mode

[M. Jia *et al.*, *NF* 61, 106023(2021)]

- Advantages of $n = 4$ RMPs
 - No obvious drop of energy confinement
 - Avoidance of W accumulation

Enhanced core confinement via triggering ITB during the application of $n=4$ RMPs






- ITB was triggered reproducibly during the application of $n=4$ RMP ELM control
- ITB can be stably sustained by RMP




Simultaneously control of double transport barriers with low core W

RMP ELM suppression strongly depends on the modification of pedestal profiles



- Reduced pedestal gradient due to **increase of $n_{e,sep}$** during the transition from ELM mitigation to **suppression**
- Peeling-ballooning modes are stabilized by the reduction of pedestal current and pressure gradient induced by RMPs

Resonant component plasma response plays the dominant role in achieving ELM suppression






[Sun Y. *et al*, FEC(2025)]

- Resonant response in $[q_{95}, n_e]$ window aligns well with ELM suppression

- Resonant RMP on ELM suppression
- Non-resonant RMP on fast-ion loss
- Optimized RMP ELM suppression with minimal fast-ion loss

RMP effects on AE and fast particle loss






[Chu et al, NF 58, 104004(2018)] [Garcia-Munoz et al, PPCF 61, 054007(2019)] [He et al, NF 59, 126026 (2019)]

- RMP causes the changes of the stability of AE
- Various resonant effects have been identified



Phase space profile control

Non-resonant RMP side bands enhanced fast-ion loss





[He K. *et al*, NF 60, 126027 (2020)]

- Plasma response can significantly shield the resonant components of RMPs
- RMPs with plasma response enhanced EP loss due to side band resonances

Far non-resonant RMP components effect on fast-ion loss during the application of $n=4$ RMPs

Drift islands
($n=4$):

$$m_p = 9$$



$$\hat{B}_{m_p} = \frac{1}{2\pi} \int_0^{2\pi} e^{-im_p\theta_p} \left[\sum_{m_b} B_{m_b}(\psi) e^{i(m_b\theta - n\Delta)} \right] d\theta_p$$

In orbit coordinates




In flux coordinates

[Zhang YN *et al*, NF 64, 046012
(2024)]

- Significant fast-ion loss even with strong shielding effect


Why far non-resonant harmonics matter

Experienced field: $\sum_{m_b} B_{m_b}(\psi) e^{i(m_b\theta - n\Delta)}$




- Edge non-resonant RMPs resonant with fast-ions with finite orbit width and variation of toroidal drift
- RFA effects due to kink mode response further enhance the non-resonant harmonics, although there is a shielding of resonant

Magnetic topology .VS. particle orbit topology





Magnetic islands

$$w_{mn} = 4 \sqrt{\left| \frac{b_{mn}}{nq'/q} \right|}_{q=m/n}$$



Drift islands

$$\Delta P_\zeta = 4 \sqrt{\frac{2n h_{m'n}}{\partial \dot{Q}(\mu, P_\zeta, E) / \partial P_\zeta}}$$




[He K. *et al*, NF 65, 106022 (2025)] Spectrum in orbit coordinates

$$h_{m'n}(\mu, P_\zeta, J)$$

- Non-resonant RMP contributes to the drift islands
- Drift island width can be evaluated using an action-angle



- Resonant RMP on ELM suppression
- Non-resonant RMP on fast-ion loss
- **Optimized RMP ELM suppression with minimal fast-ion loss**

Summary of physics in different scales





- Helicity of field line is different from helicity of particle orbits

Optimization of $n=2$ RMP spectrum for ELM control with minimal fast-ion loss





- Separation of phasing peaks with strongest resonant and non-resonant harmonics allows an optimization
- The **reduction** of fast-ion can be a **factor 2-4** for the same edge resonance level for RMP ELM suppression

The optimization is applicable for different operational scenarios




- The phasing difference $\Delta(\delta\phi_{UL})$ between maximal edge resonance and fast-ion loss **keeps almost constant** in a large range of q_{95}
- Optimization becomes more challenging in **advanced scenario** due to RFA




- Resonant tearing layer response is sensitive to layer quantities, such as rotation and resistivity

Another advantage of $n=4$ RMPs for ELM suppression



- $n=4$ RMPs is naturally optimized for the phasing with maximal edge resonance and minimal fast-ion loss
- Minimal level of fast-ion loss for the $n=4$ case is **smaller** than the $n=2$ case for the same level of edge resonance



Summary and conclusion

- RMP ELM suppression window observed in the EAST experiments can be well understood by the **resonant tearing-like layer physics** plasma response
- **Non-resonant kink-like** plasma response dominates the fast-ion loss in the simulation
- **Optimized RMP spectrum** for ELM suppression with minimal fast-ion loss has been identified, utilizing the different plasma response in resonant and non-resonant harmonics
- Advantages of $n=4$ RMPs for ELM suppression in both energy confinement and fast-ion loss has been identified

